| | DELAWARE MONTHLY HIV/AIDS REPORT Delaware data through 08/31/2015 | | | | | | | | | | | | |
|--|--|---------------|----------------|-------------|--------------------|---|-----------------------------|----------|-----------------|---------|---------------------|----------------|--|
| | Living ¹ w/ HIV | | Living w/ AIDS | | Living w/ HIV/AIDS | | Cumulative ² HIV | | Cumulative AIDS | | Cumulative HIV/AIDS | | |
| Exposure Category | Number | Percent | Number | | Number | Percent | Number | Percent | Number | Percent | Number | Percent | |
| Gender | | - | | | | | | | | | | | |
| Vale | 924 | 69.5% | 1499 | 68.5% | 2423 | 68.9% | 974 | 69.0% | 3189 | 72.2% | 4163 | 71.4% | |
| Female | 406 | | 688 | 31.5% | 1094 | | 437 | | 1228 | | 1665 | | |
| Total | 1330 | | 2187 | 100% | 3517 | | 1411 | 100% | 4417 | 100% | 5828 | | |
| Residence | | | | | | | | | | | | | |
| Kent County | 168 | 12.6% | 304 | 13.9% | 472 | 13.4% | 157 | 11.1% | 476 | 10.8% | 633 | 10.9% | |
| New Castle County | 887 | | 1444 | 66.0% | 2331 | | 1007 | | 3304 | | 4311 | | |
| Wilmington Metro Area | 514 | 38.6% | | 42.3% | 1440 | 40.9% | 596 | 42.2% | 2176 | 49.3% | 2772 | 47.6% | |
| Outside Wilmington Metro Area | 373 | 28.0% | | 23.7% | 891 | 25.3% | 411 | 29.1% | 1128 | 25.5% | 1539 | 26.4% | |
| Sussex County | 268 | | 430 | 19.7% | 698 | | 247 | | 637 | 14.4% | 884 | | |
| Unknown | 7 | | 9 | 0.4% | 16 | | 247 | | 007 | | 004 | | |
| Total | 1330 | | 2187 | 100% | 3517 | | 1411 | 100% | 4417 | | 5828 | | |
| Age at Diagnoses | 1000 | 10070 | 2107 | 10070 | 0011 | 10070 | | 10070 | | 10070 | 0020 | 10070 | |
| | 40 | 0.00/ | 40 | 0.00(| _ | | 10 | 0.70/ | 00 | 0.00/ | _ | | |
| <13 | 12 | | 18 | 0.8% | HIV ar | nd AIDS | 10 | | 28 | | | | |
| 13-19 | 92 | | 23 | 1.1% | diagnos | ses ages | 90 | | 30 | | | IDS diagnoses | |
| 20-29 | 404 | | 336 | 15.4% | | two different | 404 | | 588 | | • | present two | |
| 30-39 | 407 | 30.6% | 757 | 34.6% | disease : | states and | 422 | | 1654 | | | lisease states | |
| 40-49 | 280 | | 719 | 32.9% | cannot be | e combined | 320 | | 1422 | | and canno | t be combined | |
| 50+ | 135 | | 334 | 15.3% | | | 165 | | 695 | | | | |
| Total | 1330 | 100% | 2187 | 100% | _ | | 1411 | 100% | 4417 | 100% | | | |
| Race/Ethnicity | | | | | | | | | | | | | |
| Hispanic - All races | 99 | | 145 | 6.6% | 244 | | 96 | | 241 | 5.5% | 337 | | |
| White | 427 | | 639 | 29.2% | 1066 | | 427 | | 1197 | | 1624 | | |
| Black | 774 | | 1359 | 62.1% | 2133 | 60.6% | 864 | | 2936 | | 3800 | | |
| Other ³ | 30 | | 44 | 2.0% | 74 | | 24 | | 43 | | 67 | | |
| Total | 1330 | 100% | 2187 | 100% | 3517 | 100% | 1411 | 100% | 4417 | 100% | 5828 | 100% | |
| Mode of Exposure in Adult Cases | | | | | | | | | | | | | |
| MSM | 540 | 40.6% | 729 | 33.3% | 1269 | 36.1% | 531 | 37.6% | 1363 | 30.9% | 1894 | 32.5% | |
| IDU | 194 | 14.6% | 510 | 23.3% | 704 | 20.0% | 264 | 18.7% | 1559 | 35.3% | 1823 | 31.3% | |
| MSM/IDU | 33 | 2.5% | 110 | 5.0% | 143 | 4.1% | 44 | 3.1% | 232 | 5.3% | 276 | 4.7% | |
| Heterosexual contact w/IDU | 49 | 3.7% | 160 | 7.3% | 209 | 5.9% | 56 | 4.0% | 362 | 8.2% | 418 | 7.2% | |
| Heterosexual contact | 411 | 30.9% | 569 | 26.0% | 980 | 27.9% | 437 | 31.0% | 743 | 16.8% | 1180 | 20.2% | |
| Transfusion/transplant recipient | 1 | 0.1% | 4 | 0.2% | 5 | 0.1% | 1 | 0.1% | 18 | 0.4% | 19 | 0.3% | |
| Risk not reported/other | 90 | 6.8% | 70 | 3.2% | 160 | 4.5% | 68 | 4.8% | 94 | 2.1% | 162 | 2.8% | |
| Pediatric Exposure | 12 | 0.9% | 35 | 1.6% | 47 | 1.3% | 10 | 0.7% | 46 | 1.0% | 56 | 1.0% | |
| Total | 1330 | 100% | 2187 | 100% | 3517 | 100% | 1411 | 100% | 4417 | 100% | 5828 | 100% | |
| wode of Exposure in Pediatric Cases | | | | | | | | | | | | | |
| Mother with/at risk for infection | 9 | 75.0% | 28 | 80.0% | 37 | 78.7% | 8 | 80.0% | 41 | 89.1% | 49 | 87.5% | |
| Receipt of blood transfusion or tissue | 0 | | 1 | 2.9% | 1 | | 0 | | 1 | 2.2% | 1 | 1.8% | |
| Hemophillia/Coagulation Disorder | 0 | | 0 | 0.0% | 0 | | 0 | | 1 | 2.2% | 1 | 1.8% | |
| Other/risk not reported | 3 | | 6 | 17.1% | g | | 2 | | 3 | | 5 | | |
| Total | 12 | | 35 | 100% | 47 | | 10 | | 46 | | 56 | | |
| "Living" data includes anyone currently living in DE with HIV/AIDS. | | | | | | Eight Year History of New Infections (DE Cases) | | | | | | | |
| ² Cumulative data includes only those people who were residents of DE at time of diagnosis. | | | | | | | | | | | | | |
| ³ Other includes Asian, Pacific Islander, American Indian, Alaskan Native or Multi-racial. | | | | | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | |
| | | | | | 160 | 153 | 131 | 109 | 136 | 115 | 117 | 65 | |
| Pediatric cases are individuals <13 years of age. | | | | | | | *So | Far This | Year | | | | |
| 14.89% of HIV/AIDS diagnosed persons livir | ig in DE we | re diagnose | d in another | state and r | noved to DE | after Diagnos | ses | | | | | _ | |
| e . | - | IV cases sinc | | | | | ar in whic | | | e | | | |