



DELAWARE MONTHLY HIV/AIDS REPORT

Delaware data through 12/31/2016

| Exposure Category | Living ¹ w/ HIV | | Living w/ AIDS | | Living w/ HIV/AIDS | | Cumulative ² HIV | | Cumulative AIDS | | Cumulative HIV/AIDS | |
|--|----------------------------|-------------|----------------|-------------|---|-------------|-----------------------------|-------------|-----------------|--|---------------------|-------------|
| | Number | Percent | Number | Percent | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| Gender | | | | | | | | | | | | |
| Male | 923 | 70.9% | 1427 | 68.6% | 2350 | 69.5% | 1029 | 69.6% | 3258 | 72.4% | 4287 | 71.7% |
| Female | 379 | 29.1% | 652 | 31.4% | 1031 | 30.5% | 450 | 30.4% | 1244 | 27.6% | 1694 | 28.3% |
| Total | 1302 | 100% | 2079 | 100% | 3381 | 100% | 1479 | 100% | 4502 | 100% | 5981 | 100% |
| Residence | | | | | | | | | | | | |
| Kent County | 169 | 13.0% | 284 | 13.7% | 453 | 13.4% | 170 | 11.5% | 488 | 10.8% | 658 | 11.0% |
| New Castle County | 853 | 65.5% | 1373 | 66.0% | 2226 | 65.8% | 1048 | 70.9% | 3363 | 74.7% | 4411 | 73.8% |
| Wilmington Metro Area | 468 | 35.9% | 859 | 41.3% | 1327 | 39.2% | 614 | 41.5% | 2208 | 49.0% | 2822 | 47.2% |
| Outside Wilmington Metro Area | 385 | 29.6% | 514 | 24.7% | 899 | 26.6% | 434 | 29.3% | 1155 | 25.7% | 1589 | 26.6% |
| Sussex County | 276 | 21.2% | 417 | 20.1% | 693 | 20.5% | 261 | 17.6% | 651 | 14.5% | 912 | 15.2% |
| Unknown | 4 | 0.3% | 5 | 0.2% | 9 | 0.3% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| Total | 1302 | 100% | 2079 | 100% | 3381 | 100% | 1479 | 100% | 4502 | 100% | 5981 | 100% |
| Age at Diagnoses | | | | | | | | | | | | |
| <13 | 13 | 1.0% | 19 | 0.9% | HIV and AIDS diagnoses ages represent two different disease states and cannot be combined | 10 | 0.7% | 28 | 0.6% | HIV and AIDS diagnoses ages represent two different disease states and cannot be combined | | |
| 13-19 | 86 | 6.6% | 25 | 1.2% | | 92 | 6.2% | 31 | 0.7% | | | |
| 20-29 | 410 | 31.5% | 313 | 15.1% | | 436 | 29.5% | 605 | 13.4% | | | |
| 30-39 | 391 | 30.0% | 721 | 34.7% | | 437 | 29.5% | 1678 | 37.3% | | | |
| 40-49 | 265 | 20.4% | 677 | 32.6% | | 326 | 22.0% | 1440 | 32.0% | | | |
| 50+ | 137 | 10.5% | 324 | 15.6% | | 178 | 12.0% | 720 | 16.0% | | | |
| Total | 1302 | 100% | 2079 | 100% | 1479 | 100% | 4502 | 100% | | | | |
| Race/Ethnicity | | | | | | | | | | | | |
| Hispanic - All races | 94 | 7.2% | 132 | 6.3% | 226 | 6.7% | 98 | 6.6% | 251 | 5.6% | 349 | 5.8% |
| White | 433 | 33.3% | 620 | 29.8% | 1053 | 31.1% | 445 | 30.1% | 1217 | 27.0% | 1662 | 27.8% |
| Black | 740 | 56.8% | 1277 | 61.4% | 2017 | 59.7% | 909 | 61.5% | 2990 | 66.4% | 3899 | 65.2% |
| Other ³ | 35 | 2.7% | 50 | 2.4% | 85 | 2.5% | 27 | 1.8% | 44 | 1.0% | 71 | 1.2% |
| Total | 1302 | 100% | 2079 | 100% | 3381 | 100% | 1479 | 100% | 4502 | 100% | 5981 | 100% |
| Mode of Exposure in Adult Cases | | | | | | | | | | | | |
| MSM | 554 | 42.5% | 703 | 33.8% | 1257 | 37.2% | 566 | 38.3% | 1398 | 31.1% | 1964 | 32.8% |
| IDU | 158 | 12.1% | 458 | 22.0% | 616 | 18.2% | 263 | 17.8% | 1563 | 34.7% | 1826 | 30.5% |
| MSM/IDU | 35 | 2.7% | 94 | 4.5% | 129 | 3.8% | 43 | 2.9% | 234 | 5.2% | 277 | 4.6% |
| Heterosexual contact w/IDU | 44 | 3.4% | 157 | 7.6% | 201 | 5.9% | 54 | 3.7% | 367 | 8.2% | 421 | 7.0% |
| Heterosexual contact | 399 | 30.6% | 548 | 26.4% | 947 | 28.0% | 458 | 31.0% | 770 | 17.1% | 1228 | 20.5% |
| Transfusion/transplant recipient | 1 | 0.1% | 4 | 0.2% | 5 | 0.1% | 1 | 0.1% | 18 | 0.4% | 19 | 0.3% |
| Risk not reported/other | 98 | 7.5% | 80 | 3.8% | 178 | 5.3% | 84 | 5.7% | 106 | 2.4% | 190 | 3.2% |
| Pediatric Exposure | 13 | 1.0% | 35 | 1.7% | 48 | 1.4% | 10 | 0.7% | 46 | 1.0% | 56 | 0.9% |
| Total | 1302 | 100% | 2079 | 100% | 3381 | 100% | 1479 | 100% | 4502 | 100% | 5981 | 100% |
| Mode of Exposure in Pediatric Cases | | | | | | | | | | | | |
| Mother with/at risk for infection | 10 | 76.9% | 31 | 88.6% | 41 | 85.4% | 8 | 80.0% | 41 | 89.1% | 49 | 87.5% |
| Receipt of blood transfusion or tissue | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 1 | 2.2% | 1 | 1.8% |
| Hemophilia/Coagulation Disorder | 0 | 0.0% | 1 | 2.9% | 1 | 2.1% | 0 | 0.0% | 1 | 2.2% | 1 | 1.8% |
| Other/risk not reported | 3 | 23.1% | 3 | 8.6% | 6 | 12.5% | 2 | 20.0% | 3 | 6.5% | 5 | 8.9% |
| Total | 13 | 100% | 35 | 100% | 48 | 100% | 10 | 100% | 46 | 100% | 56 | 100% |

¹ "Living" data includes anyone currently living in DE with HIV/AIDS.

² Cumulative data includes only those people who were residents of DE at time of diagnosis.

³ Other includes Asian, Pacific Islander, American Indian, Alaskan Native or Multi-racial.

^{*} Pediatric cases are individuals <13 years of age.

Eight Year History of New Infections (DE Cases)

| 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 |
|-------------------|------|------|------|------|------|------|------|
| 131 | 109 | 138 | 113 | 117 | 101 | 105 | 0 |
| *So Far This Year | | | | | | | |

14.89% of HIV/AIDS diagnosed persons living in DE were diagnosed in another state and moved to DE after Diagnoses

Note: AIDS cases have been reported in Delaware since 1983, HIV cases since July 2001.

New Infection = Year in which HIV infection was first diagnosed